

# Clinimix

## **To open:**

Tear overwrap across top at slit and remove solution container. Some opacity (fogginess) of the plastic due to moisture absorption during the sterilization process may be observed, and is normal. This does not affect the solution quality or safety, and will gradually fade.

Check to ensure the seal between the chambers is intact, and that the solutions are contained in separate chambers. Check for small leaks by separately squeezing each chamber. Notify pharmacy if either of these are observed, as sterility or stability of the solution may be affected.

## **To Activate:**

Be sure to activate the bag and thoroughly mix the amino acid (protein) and dextrose (sugar) components prior to adding any additives. Large volume additives should be transferred into the bag prior to smaller volume additives.



Lay the bag onto a flat surface. Grasp the bag firmly with both hands at the top corners.



Starting from the top, squeeze and roll the bag down toward the bottom until the peel-seal is broken down the center.



Shown: CLINDIMIX Injections bag

You should feel or hear an audible pop as the peel-seal dividing the chambers is broken, which allows the components from each chamber to begin to mix.



Shown: CLINDIMIX Injections bag

It's important to make sure the peel-seal is broken the entire length of the bag, as shown in the bag on the right where the chambers are no longer separated.

If the seal has not been separated completely, flip the bag over with the port side down and roll the opposite direction until the seal fully separates. **Do not pull the bag apart to separate the seal.**



Shown: CLINDIMIX Injections bag

Hold the bag up and squeeze from side to side to thoroughly mix the solutions. Check for leaks. If leaks are found, discard the product as sterility or stability may be impaired.

## To Add Medications (example Infuvite):



Prepare Injection port by swabbing with an alcohol prep pad using a back and forth motion. Inject the ordered additives through the medication injection port. Mix solution and medications thoroughly. Check for leaks.

## Administration



Twist off protector cap from administration port. Attach TPN administration set by spiking bag in administration port **using a twisting motion**. Attach to infusion pump.

## Storage:

Keep Clinimix in protective overwrap at room temperature until ready for use. Unmixed solution is stable through the printed date on package. Once protective overwrap is removed and solution is mixed, it is stable in the refrigerator for up to 9 days. Additives to Clinimix may not be stable for more than 24 hours-check with pharmacy for specific information.